



ELSEVIER

Contents lists available at ScienceDirect

Biochemical Systematics and Ecology

journal homepage: www.elsevier.com/locate/biochemsyseco

Volatile components from the mandibular glands of the turtle ants, *Cephalotes alfaroi* and *Cephalotes cristatus*

William F. Wood^{a,*}, Thuy-Tien Hoang^b, Terrence P. McGlynn^b^a Department of Chemistry, Humboldt State University, Arcata, CA 95521, USA^b Department of Biology, California State University, Dominguez Hills, Carson, CA 90747, USA

ARTICLE INFO

Article history:

Received 14 December 2010

Accepted 29 January 2011

Available online xxx

Keywords:

*Cephalotes alfaroi**Cephalotes cristatus*

Hymenoptera

Formicidae

Myrmicinae

Cephalotini

Mandibular gland secretion

Turtle ant

4-Heptanone

4-Heptanol

ABSTRACT

GC–MS analysis of whole head extracts from the turtle ants, *Cephalotes alfaroi* and *Cephalotes cristatus*, showed that 4-heptanone and 4-heptanol were the major volatile components in the mandibular glands. 4-Heptanone and 4-heptanol have rarely been identified in mandibular gland secretions from other ant genera. Thus, these compounds may be chemotaxonomic markers for the genus *Cephalotes*, since they have been identified in the mandibular glands from all members of this genus that have been investigated to date. Minor components identified in the whole head extracts of these ants were 3-methyl-1-butanol, 3-heptanone, 3-hexanol, 2- and 3-methylbutanoic acids, 2-methyl-4-heptanone, 2-phenylethanol and phenol. To our knowledge, this is the first time that 2-methyl-4-heptanone and phenol have been reported in the mandibular gland secretion from any Formicid.

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

Ant mandibular gland secretions were among the first semiochemicals to be identified. Over the years, many different chemicals have been documented from these glands (Ali et al., 1987; Attygalle and Morgan, 1984; Bellas and Hölldobler, 1985; Bestmann et al., 1988; Blum, 1969, 1981; Blum and Hermann, 1978a,b; Bradshaw et al., 1979; Cammaerts et al., 1983, 1987; do Nascimento et al., 1993; Hölldobler and Wilson, 1990; Hughes et al., 2001; Jackson and Morgan, 1993; Jones et al., 2004; Keegans et al., 1991, 1992; Morgan et al., 1999, 2006, 2008; Oldham and Morgan, 1993; Oldham et al., 1994; Olubajo et al., 1980; Parry and Morgan, 1979; Voegtli et al., 2008; Wood, 2005; Wood and Chong, 1975; Wood et al., 2002, 2006). Ketones and their corresponding alcohols are reported as the major components in many of these secretions. In almost all cases, the carbonyl or corresponding hydroxy group of these compounds is either at the 2- or 3-position of the carbon chains. There are few reports of ketones and alcohols at the 4-position of the carbon chains from ant mandibular gland secretions.

4-Heptanone and 4-heptanol were identified in the mandibular gland extracts of the turtle ants (Myrmicinae), *Cephalotes varians* and *Cephalotes pusillus* (Olubajo et al., 1980; Morgan et al., 2006). The only other ant species reported to contain both 4-heptanone and 4-heptanol in its mandibular glands is the unrelated African Formicine, *Polyrhachis schistacea* (Brand and Lindner, 1989). Finally, 4-heptanone along with 6-methyl-5-hepten-2-one and 6-methyl-5-hepten-2-ol has been reported from the mandibular glands of the weaver ant (Formicinae) *Polyrhachis simplex* (Hefetz and Lloyd, 1982).

* Corresponding author. Tel.: +1 707 826 3109; fax: +1 707 826 3279.

E-mail address: wfw2@humboldt.edu (W.F. Wood).

Ants of the genus *Cephalotes* are large, attractive, heavily armoured and arboreal residents of the high rain forest canopies in the Neotropics. Because of the inaccessible locations that many members of this genus inhabit, it is not surprising that only two previously noted members of the genus *Cephalotes* have been investigated to date (Olubajo et al., 1980; Morgan et al., 2006). We examined whole head extracts from two additional members of this genus, *Cephalotes alfaroi* and *Cephalotes cristatus*. Like the previous investigations on members of this genus, we found that the major components in the mandibular gland secretions of *C. alfaroi* and *C. cristatus* were 4-heptanone and 4-heptanol. Unlike previous reports on *Cephalotes*' mandibular glands, we report minor compounds in the whole head extracts from *C. alfaroi* and *C. cristatus*.

2. Materials and methods

Workers of the arboreal turtle ants, *C. alfaroi* and *C. cristatus* were collected at La Selva Biological Station, Heredia Province, Costa Rica (10° 25' 54" N, 84° 00' 13" W) and were identified by TM using online taxonomic resources (Longino, 2009). *C. alfaroi* workers were individually collected from trees, placed in resealable plastic bags, and then put in a freezer for over 1 h to immobilise the ants. The heads from 25 worker ants were excised and placed in glass vials with Teflon-lined stoppers containing 0.5 mL of dichloromethane (CH₂Cl₂), which was stored at –20 °C for later analysis. Likewise, two additional collections of 25 *C. alfaroi* worker ant heads were made (the three collections were analyzed separately and the results averaged.) Finally, a single collection of 30 *C. cristatus* worker ant heads was made using the same procedure. The CH₂Cl₂ from these extractions was analyzed without any further manipulation. Decapitation and storage of ant heads in solvent is a standard method for the extraction of the mandibular gland secretions of ants when they cannot be analyzed immediately (Wood, 2005; Wood et al., 2002, 2006). A recent study showed no difference between whole head extracts and pure mandibular secretions of the congeneric *C. pusillus* (Morgan et al., 2006).

GC–MS analyses were performed on the CH₂Cl₂ extracts in a splitless mode (0.5 min), using a Hewlett–Packard GCD Plus fitted with a 30 m × 0.25 μm cross-linked phenyl methyl silicone capillary column (HP-5MS). The gas chromatograph was programmed so the oven temperature was kept at 40 °C for 4 min, then increased to a final temperature of 325 °C at a rate of 30 °C/min and held at this temperature for 5 min. Impurities in control solvent samples and sample constituents less than 0.1% of the total ion current (TIC) were not reported. Initial identification of secretion components was done by comparison to published electron impact - mass spectra (EI-MS) in the National Institute of Standards and Technology (NIST) 1998 computerised mass spectral library, and confirmed by comparison of spectra and retention times to those of commercially available standards, except 2-methyl-4-heptanone that was prepared synthetically (de Jong and Feringa, 1991).

3. Results

The major compounds in the mandibular glands of *C. alfaroi* and *C. cristatus* were 4-heptanone and 4-heptanol (Table 1). Previous reports of 4-heptanone and 4-heptanol in mandibular glands of *C. varians* and *C. pusillus* did not identify any additional components (Olubajo et al., 1980; Morgan et al., 2006). In our study, we found 3-methyl-1-butanol, 3-heptanone, 3-hexanol, 2- and 3-methylbutanoic acids, 2-methyl-4-heptanone, phenol, and 2-phenylethanol as minor components in the whole head extracts (Table 1).

4. Discussion and conclusion

Ant mandibular gland secretions often function as alarm pheromones, as they typically elicit aggressive behaviour. Ants of the genus *Cephalotes* are generally unaggressive and frequently use a passive defence that includes stopping and lying flat on the surface when molested. Several studies have reported on the defensive behaviour of *C. varians*, a species that ranges from the Florida Keys to the Bahamas, Cuba and Jamaica. Wilson (1976) reported an ethogram, a complete catalogue of the behavioural patterns of this species. Another study describes the behaviour of *C. varians* workers in response to mandibular

Table 1
Volatile compounds in *Cephalotes* mandibular glands as % of the total ion current (TIC).

Compound	Retention time (min)	<i>C. alfaroi</i> ^a	<i>C. cristatus</i> ^b
3-methyl-1-butanol	3.45	–	0.7
3-Hexanone	4.54	1.5	0.1
3-Hexanol	4.70	1.5	0.1
3-Methylbutanoic acid	5.69	0.4	–
2-Methylbutanoic acid	5.74	0.4	–
4-Heptanone	5.86	87.4	49.3
4-Heptanol	5.99	6.3	46.9
2-Methyl-4-heptanone	6.34	–	0.3
Phenol	6.86	2.6	1.9
2-Phenylethanol	7.97	–	0.7

^a Average of 3 replicate analyses of pooled extracts from 25 workers.

^b Single sample analysis of pooled extracts from 30 worker.

gland chemicals (Olubajo et al., 1980). When worker ants were exposed either to crushed ant heads, 4-heptanone, 4-heptanol, or a mixture of these chemicals, a non-aggressive response was elicited. The ants were reported to stop moving and back up when exposed to these chemical cues. 4-Heptanone has been tested for alarm behaviour with *Pogonomyrmex badius* (Blum et al., 1971) and *Atta texana* (Moser et al., 1968), ants that utilise 4-methyl-3-heptanone as an alarm pheromone. In both cases, 4-heptanone was weakly active and only elicited alarm behaviour at high concentrations. Thus, it is possible that ant species, which use alarm pheromones with a ketone functional group in the 2- or 3-position, are unable to detect chemical cues from *Cephalotes* spp.

A major difference between 4-heptanol and mandibular gland alcohols with the hydroxy group in the 2- or 3-position, is that it is achiral. Mandibular gland alcohols with the hydroxy group in the 2- or 3-position are chiral, and it has been shown that ants produce enantiomerically pure compounds. As an example, *Crematogaster castanea* and *Crematogaster liengmei* make only (S)-(+)-3-octanol (Brand, 1985) and (S)-2-methyl-1-hexanol is the only isomer produced by *Cataglyphis bicolor* (Agosti et al., 1996). The lack of chirality may be important to the semiochemistry of 4-heptanol.

The minor compounds, except phenol and 2-methyl-4-heptanone, have frequently been reported in ant mandibular gland secretions. 2-Methyl-4-heptanone has been reported from the anal (pygidial) glands of *Tapinoma simrothi* (Hefetz and Lloyd, 1983) and *Tapinoma nigerrimum* (Trave and Pavan, 1956). Phenol has been reported from the Dufour's gland of several *Myrmecocystus* spp. (Lloyd et al., 1989) and the poison gland of *Camponotus socius* (Kohl et al., 2001). To our knowledge, this is the first time that 2-methyl-4-heptanone and phenol have been reported in the mandibular gland secretion from any Formicid.

Finally, if it is found that other members of genus *Cephalotes* also have high concentrations of 4-heptanone and 4-heptanol in their mandibular gland secretions, these compounds may be chemotaxonomic markers for the genus.

Acknowledgements

This project was supported by the U.S. National Science Foundation (OISE-0749047; OISE-0854259) and a CSUDH RSCAAP award to TM.

References

- Agosti, D., Austin, C., Gökçen, O.A., König, W.A., Morgan, E.D., Scott, E.D., Wehner, R., 1996. (S)-2-methyl-1-hexanol, a characteristic mandibular gland substance of ants of the *Cataglyphis bicolor* group. *Chemoecology* 7, 57–60.
- Ali, M.F., Cammaerts, R., Evershed, R.P., Morgan, E.D., 1987. A reexamination of the mandibular gland contents of *Tetramorium caespitum* (Hymenoptera: Formicidae). *Insect Biochem.* 17, 237–241.
- Attygalle, A.B., Morgan, E.D., 1984. Chemicals from the glands of ants. *Chem. Soc. Rev.* 13, 245–278.
- Bellas, T., Hölldobler, B., 1985. Constituents of mandibular and Dufour's glands of an Australian *Polyrhachis* weaver ant. *J. Chem. Ecol.* 11, 525–538.
- Bestmann, H.J., Attygalle, A.B., Glasbrenner, J., Riemer, R., Vostrowsky, O., Constantino, M.G., Melikyan, G., Morgan, E.D., 1988. Identification of the volatile components of the mandibular gland secretion of the ant *Manica rubida*: structure elucidation, synthesis, and absolute configuration of manicone. *Lieb. Ann. der Chem.* 1988, 55–60.
- Blum, M.S., 1969. Alarm pheromones. *Ann. Rev. Ent.* 14, 57–80.
- Blum, M.S., 1981. Chemical Defenses of Arthropods. Academic Press, New York. 452pp.
- Blum, M.S., Hermann, H.R., 1978a. Venoms and venom apparatuses of the Formicidae: Myrmecinae, Ponerinae, Dorylinae, Pseudomyrmecinae, Myrmicinae, and Formicinae. In: Bertini, S. (Ed.), *Arthropod Venoms. Handb. Exp. Pharm.* 48, pp. 870–894.
- Blum, M.S., Hermann, H.R., 1978b. Venoms and venom apparatuses of the Formicidae: Dolichoderinae and Aneuretinae. In: Bertini, S. (Ed.), *Arthropod Venoms. Handb. Exp. Pharm.* 48, pp. 801–869.
- Blum, M.S., Doolittle, R.E., Beroza, M., 1971. Alarm pheromones: utilization in evaluation of olfactory theories. *J. Insect Physiol.* 17, 2351–2361.
- Bradshaw, J.W.S., Baker, R., Howse, P.E., 1979. Multicomponent alarm pheromones in the mandibular glands of major workers of the African weaver ant, *Oecophylla longinoda*. *Physiol. Ent.* 4, 15–25.
- Brand, J.M., 1985. Enantiomeric composition of an alarm pheromone component of the ants *Crematogaster castanea* and *C. liengmei*. *J. Chem. Ecol.* 11, 177–180.
- Brand, J.M., Lindner, W.A., 1989. Volatile constituents of a Southern African ant, *Polyrhachis schistacea* (Gerstaecker). *So. Afr. J. Sci.* 85, 388–389.
- Cammaerts, M.C., Evershed, R.P., Morgan, E.D., 1983. The volatile components of the mandibular gland secretion of workers of the ants *Myrmica lobicornis* and *Myrmica sulcinodis*. *J. Insect Physiol.* 29, 659–664.
- Cammaerts, M.C., Inwood, M.R., Morgan, E.D., Parry, K., Tyler, R.C., 1987. Comparative study of the pheromones emitted by workers of the ants *Myrmica rubra* and *Myrmica scabrinodis*. *J. Insect Physiol.* 24, 207–214.
- de Jong, E.A., Feringa, B.L., 1991. The Synthesis of 2-methyl-4-heptanone. *J. Chem. Ed.* 68, 71–72.
- do Nascimento, R.R., Jackson, B.D., Morgan, E.D., Clark, W.H., Blom, P.E., 1993. Chemical secretions of two sympatric harvester ants, *Pogonomyrmex salinus* and *Messor lobognathus*. *J. Chem. Ecol.* 19, 1993–2005.
- Hefetz, A., Lloyd, H.A., 1982. Exocrine glands of *Polyrhachis Simplex*: chemistry and function. *J. Chem. Ecol.* 8, 635–639.
- Hefetz, A., Lloyd, H.A., 1983. Identification of new components from anal glands of *Tapinoma simrothi* pheonicium. *J. Chem. Ecol.* 9, 607–613.
- Hölldobler, B., Wilson, E.O., 1990. *The Ants*. The Belknap Press of Harvard University Press, Cambridge, Mass. 732pp.
- Hughes, W.O.H., House, P.E., Goulson, D., 2001. Mandibular gland chemistry of grass-cutting ants: species, caste, and colony variation. *J. Chem. Ecol.* 27, 109–124.
- Jackson, B.D., Morgan, E.D., 1993. Insect chemical communication: pheromones and exocrine glands of ants. *Chemoecology* 4, 125–144.
- Jones, T.H., Clark, D.A., Edwards, A.A., Davidson, D.W., Spande, T.F., Snelling, R.R., 2004. The chemistry of exploding ants, *Camponotus* spp. (cylindricus COMPLEX). *J. Chem. Ecol.* 30, 1479–1492.
- Keegans, S.J., Billen, J., Morgan, E.D., 1991. Volatile secretions of the green tree ant *Oecophylla smaragdina* (Hymenoptera: Formicidae). *Comp. Biochem. Physiol.* 100B, 681–685.
- Keegans, S.J., Morgan, E.D., Agosti, D., Wehner, R., 1992. What do glands tell us about species? A chemical case study of *Cataglyphis* ants. *Biochem. Syst. Ecol.* 20, 559–572.
- Kohl, E., Hölldobler, B., Bestmann, H.J., 2001. Trail and recruitment pheromones in *Camponotus socius* (Hymenoptera: Formicidae). *Chemoecology* 11, 67–73.
- Lloyd, H.A., Blum, M.S., Snelling, R.R., Evans, S.L., 1989. Chemistry of mandibular and Dufour's gland secretions of ants in genus *Myrmecocystus*. *J. Chem. Ecol.* 15, 2589–2599.

- Longino, J.T., 2009. Ants of Costa Rica - Cephalotes Species Known from Costa Rica. <http://academic.evergreen.edu/projects/ants/genera/cephalotes/specieslist.html> (Accessed September 2009).
- Morgan, E.D., do Nascimento, R.R., Keegans, S.J., Billen, J., 1999. Comparative study of mandibular gland secretions of workers of ponerine. J. Chem. Ecol. 25, 1395–1409.
- Morgan, D.E., Keegans, S.J., Billens, J., 2006. Exocrinechemistry of the Myrmicine ant *Zacryptocerus pusillus* (Hymenoptera: Formicidae). Arch. Insect Biochem. Physiol. 63, 11–14.
- Morgan, E.D., Jungnickel, H., Billen, J., Ito, F., Bergmann, J., Gobin, B., 2008. Contents of the exocrine glands of the ant subfamily Cerapachyinae. Biochem. Syst. Ecol. 36, 260–265.
- Moser, J.C., Brownlee, R.C., Silverstein, R., 1968. Alarm pheromones of the ant *Atta texana*. J. Insect Physiol. 14, 529–535.
- Oldham, N.J., Keegans, S.J., Morgan, E.D., Paiva, R.V.S., Brandao, C.R.F., Schoeters, E., Billen, J., 1994. Mandibular gland contents of a colony of the queenless ponerine ant *Dinoponera australis*. Naturwissenschaften 81, 313–316.
- Oldham, N.J., Morgan, E.D., 1993. Structures of the pyrazines from the mandibular gland secretion of the ponerine ant *Dinoponera australis*. J. Chem. Soc. Perkin Trans., 2713–2716.
- Olubajo, O., Duffield, R.M., Wheeler, J.W., 1980. 4-Heptanone in the mandibular gland secretion of the nearctic ant, *Zaeryptoerues varians* (Hymenoptera: Formicidae). Ann. Entomol. Soc. Am. 73, 93–94.
- Parry, K., Morgan, E.D., 1979. Pheromones of ants: a review. Physiol. Entomol. 4, 161–189.
- Trave, R., Pavan, M., 1956. Veleni degli insetti. Principi estratti dalla formica *Tapinoma nigerrimum* Nyl. Chem. Ind. Milano 38, 1015–1019.
- Voegtli, H.L., Jones, T.H., Davidson, D.W., Snelling, R.R., 2008. E-2-Ethylhexenal, E-2-Ethyl-2-Hexenol, Mellein, and 4-Hydroxymellein in *Camponotus* species from Brunei. J. Chem. Ecol. 34, 215–219.
- Wilson, E.O., 1976. A social ethogram of the neotropical arboreal ant *Zacryptocerus varians* (Fr. Smith). Anim. Behav. 24, 354–363.
- Wood, W.F., 2005. Comparison of mandibular gland volatiles from ants of the bull horn acacia, *Acacia collinsii*. Biochem. Syst. Ecol. 33, 651–658.
- Wood, W.F., Chong, B., 1975. 3-Octanone and 3-octanol; alarm pheromones from East African acacia ants. J. Ga. Entomol. Soc. 10, 332–334.
- Wood, W.F., Palmer, T.M., Stanton, M.L., 2002. A comparison of volatiles in mandibular glands from three *Crematogaster* ant symbionts of the whistling thorn acacia. Biochem. Syst. Ecol. 30, 217–222.
- Wood, W.F., Palmer, T.M., Stanton, M.L., 2006. Volatiles in the mandibular gland of *Tetraoponera penzigi*: a plant ant of the whistling thorn acacia. Biochem. Syst. Ecol. 34, 536–538.